

## University of Groningen

### Moderate Traumatic Brain Injury

Einarsen, Cathrine Elisabeth; van der Naalt, Joukje; Jacobs, Bram; Follestad, Turid; Moen, Kent Gøran; Vik, Anne; Håberg, Asta Kristine; Skandsen, Toril

*Published in:*  
World neurosurgery

*DOI:*  
[10.1016/j.wneu.2018.03.176](https://doi.org/10.1016/j.wneu.2018.03.176)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2018

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Einarsen, C. E., van der Naalt, J., Jacobs, B., Follestad, T., Moen, K. G., Vik, A., Håberg, A. K., & Skandsen, T. (2018). Moderate Traumatic Brain Injury: Clinical Characteristics and a Prognostic Model of 12-Month Outcome. *World neurosurgery*, 114, e1199-e1210. <https://doi.org/10.1016/j.wneu.2018.03.176>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*



## Moderate Traumatic Brain Injury: Clinical Characteristics and a Prognostic Model of 12-Month Outcome

Cathrine Elisabeth Einarsen<sup>1,4</sup>, Joukje van der Naalt<sup>6</sup>, Bram Jacobs<sup>6</sup>, Turid Follstad<sup>5</sup>, Kent Gøran Moen<sup>4,7</sup>, Anne Vik<sup>2,4</sup>, Asta Kristine Håberg<sup>3,4</sup>, Toril Skandsen<sup>1,4</sup>

■ **BACKGROUND:** Patients with moderate traumatic brain injury (TBI) often are studied together with patients with severe TBI, even though the expected outcome of the former is better. Therefore, we aimed to describe patient characteristics and 12-month outcomes, and to develop a prognostic model based on admission data, specifically for patients with moderate TBI.

■ **METHODS:** Patients with Glasgow Coma Scale scores of 9–13 and age  $\geq 16$  years were prospectively enrolled in 2 level I trauma centers in Europe. Glasgow Outcome Scale Extended (GOSE) score was assessed at 12 months. A prognostic model predicting moderate disability or worse (GOSE score  $\leq 6$ ), as opposed to a good recovery, was fitted by penalized regression. Model performance was evaluated by area under the curve of the receiver operating characteristics curves.

■ **RESULTS:** Of the 395 enrolled patients, 81% had intracranial lesions on head computed tomography, and 71% were admitted to an intensive care unit. At 12 months, 44% were moderately disabled or worse (GOSE score  $\leq 6$ ), whereas 8% were severely disabled and 6% died (GOSE

score  $\leq 4$ ). Older age, lower Glasgow Coma Scale score, no day-of-injury alcohol intoxication, presence of a subdural hematoma, occurrence of hypoxia and/or hypotension, and preinjury disability were significant predictors of GOSE score  $\leq 6$  (area under the curve = 0.80).

■ **CONCLUSIONS:** Patients with moderate TBI exhibit characteristics of significant brain injury. Although few patients died or experienced severe disability, 44% did not experience good recovery, indicating that follow-up is needed. The model is a first step in development of prognostic models for moderate TBI that are valid across centers.

### INTRODUCTION

Few studies have specifically focused on characteristics and prognosis in patients with moderate traumatic brain injury (TBI),<sup>1–5</sup> which in existing classifications is either defined by a Glasgow Coma Scale (GCS) score of 9–12 or 9–13 at emergency department (ED) admission.<sup>6–9</sup> Previous studies of patients

#### Key words

- Cohort studies
- Comparative study
- Craniocerebral trauma
- Prognosis
- Statistical models

#### Abbreviations and Acronyms

**AUC:** Area under the curve  
**CI:** Confidence interval  
**CT:** Computed tomography  
**ED:** Emergency department  
**EDH:** Epidural hematoma  
**GCS:** Glasgow Coma Scale  
**GOSE:** Glasgow Outcome Scale Extended  
**HL test:** Hosmer-Lemeshow goodness-of-fit test  
**ICP:** Intracranial pressure  
**ICU:** Intensive care unit  
**ISS:** Injury Severity Score  
**ISSe:** Modified ISS score for extracranial injuries  
**IQR:** Interquartile range  
**SDH:** Subdural hematoma

**TBI:** Traumatic brain injury

**tSAH:** Traumatic subarachnoid hemorrhage

From the Departments of <sup>1</sup>Physical Medicine and Rehabilitation, <sup>2</sup>Neurosurgery, and <sup>3</sup>Radiology and Nuclear Medicine, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway; Departments of <sup>4</sup>Neuromedicine and Movement Science and <sup>5</sup>Public Health and Nursing, Faculty of Medicine and Health Sciences, The Norwegian University of Science and Technology, NTNU, Trondheim, Norway; <sup>6</sup>Department of Neurology AB51, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands; and <sup>7</sup>Department of Radiology, Levanger Hospital, Nord-Trøndelag Hospital Trust, Levanger, Norway

To whom correspondence should be addressed: Cathrine Elisabeth Einarsen, M.D., Ph.D. Candidate

[E-mail: [cathrine.einarsen@ntnu.no](mailto:cathrine.einarsen@ntnu.no); [cathrine.e.einarsen@gmail.com](mailto:cathrine.e.einarsen@gmail.com)]

Citation: *World Neurosurg.* (2018) 114:e1199–e1210.

<https://doi.org/10.1016/j.wneu.2018.03.176>

Journal homepage: [www.WORLDNEUROSURGERY.org](http://www.WORLDNEUROSURGERY.org)

Available online: [www.sciencedirect.com](http://www.sciencedirect.com)

1878-8750/© 2018 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

with moderate TBI have found that approximately 60% of patients had intracranial traumatic findings on admission computed tomography (CT) of the head,<sup>1-3</sup> 20%–84% were admitted to an intensive care unit (ICU),<sup>1-4</sup> and approximately 15% had surgery for a mass lesion or a depressed skull fracture.<sup>3,4</sup> However, case-fatality rates were low (0.9%–8%).<sup>2,3,5,10</sup> Furthermore, the vast majority of the patients experienced only moderate or no disability, indicating independency in daily life (74%–85%),<sup>2,5</sup> and many even a good recovery, indicating no disability (55%–75%).<sup>1,2,5</sup>

Despite the fact that the expected outcome is better after moderate than severe TBI, patients with moderate TBI are mostly studied together with patients with severe TBI in outcome prediction studies.<sup>8,11-14</sup> The largest validated prognostic models so far using the Glasgow Outcome Scale Extended (GOSE) as outcome measure are the models from the Corticosteroid randomization after significant head injury (CRASH) Trial and the International Mission for Prognosis and Analysis of Clinical Trials (IMPACT).<sup>11,15</sup> The models have consistently identified age, GCS score, pupillary reactivity, and CT characteristics as predictors for an unfavorable outcome.


These models, however, have been developed for prediction of death and severe disability (GOSE score  $\leq 4$ ), outcomes that are less likely after moderate TBI. Moreover, it has been demonstrated that these models performed better in cohorts with a high proportion of patients with poor outcomes. This was especially observed in prediction of death.<sup>16</sup> This may indicate that accurate outcome prediction in patients with better outcomes may be more challenging. Hence, there is a need for studies aiming at developing models for the prediction of outcome specifically in patients with moderate TBI, where many patients will have good recovery at follow-up. This gap in the literature also was acknowledged in a recent review.<sup>17</sup> To our knowledge, no earlier studies have constructed models for the prediction of moderate disability or worse (GOSE score  $\leq 6$ ) in contrast to a good recovery specifically in patients with moderate TBI.

Our first aim was to describe and compare clinical characteristics, head CT findings, and 12-month outcome in observational prospective cohorts of patients with moderate TBI from 2 European level I trauma centers. The second aim was to develop a prognostic model based on admission data for prediction of GOSE score  $\leq 6$ , 12 months after the injury. Prognostic models also were developed for each center separately to identify important predictors, and hence it was possible to externally validate these in the opposite dataset.

## MATERIAL AND METHODS

### The Two Centers

St. Olavs Hospital, Trondheim University Hospital (referred to as Trondheim) is a regional level I trauma center and a tertiary referral center for all neurosurgical activities for 3 counties in mid-Norway, with approximately 700,000 inhabitants and 6 general hospitals. Regarding 1 of the counties (approximately 300,000 inhabitants), patients with all severities of TBI are admitted to Trondheim. Regarding the 2 other counties, patients with severe TBI, patients with moderate TBI in need of neurosurgical




assessments and/or intervention, and patients with additional major extracranial injuries are admitted to Trondheim.

The University Medical Center in Groningen (referred to as Groningen) is 1 of 11 regional level I trauma centers in the Netherlands, serving 3 counties with 1,500,000 inhabitants in total. The area also has 12 general hospitals. Patients with all severities of TBI and patients in need for observation for neurosurgical assessments or intervention are admitted to Groningen.

### Patients and Inclusion Procedures

Patients  $\geq 16$  years of age with a moderate TBI defined by GCS score 9–13 were included. Inclusion and exclusion into the databases and follow-up in the 2 centers are described in **Figure 1A** and **B**. The GCS score was determined after stabilization in the ED. In case of intubation or sedation at the scene of accident, at



the local hospital or at the trauma center, the last nonsedated GCS score was reported ( $n = 79$ ). In Trondheim, 190 patients were enrolled prospectively in the Trondheim TBI studies during 9 years (October 2004 to September 2013) (Figure 1A). Of these patients 94, 46, and 84 have been included in previous studies from the Trondheim TBI group.<sup>4,18,19</sup> In Groningen, 205 patients were enrolled prospectively in their neurotrauma database during 8 years (January 2004 to December 2011) (Figure 1B), and 19 and 49 of these patients were included in previous studies.<sup>1,20</sup>

### Clinical Variables

Preinjury disability was defined as present if daily functioning was affected by alcohol and/or drug abuse, psychiatric or neurologic disease, developmental disorders, or severe somatic disease. Cause of injury was categorized into traffic accidents, falls, and others (including violence, ski accidents, and being struck by an object). Day-of-injury alcohol intoxication was recorded as yes or no based on the serum value of ethanol or clinical judgment, both methods have been found valid for classifying a person as sober or not.<sup>21,22</sup> Pupillary status was categorized into normal or unilaterally dilated. A secondary event was defined as occurrence of hypoxia (saturation  $<92\%$ ) and/or hypotension (systolic blood pressure  $<90$  mm Hg) at the scene of accident or at ED admission. Transfer of patients from other hospitals to the trauma centers also was recorded. Other clinical variables were being intubated, days on ventilator, treatment in a neurointensive or general intensive care unit (ICU), including the length of stay in the ICU, evacuation of any intracranial mass lesion (subdural [SDH], epidural [EDH], or intracerebral hematomas) and insertion of intracranial pressure (ICP) monitoring device (parenchymal ICP sensor and/or external ventricular drain).

The Injury Severity Score (ISS) was used to indicate overall trauma severity and was assessed by residents in neurosurgery (K.G.M. and S.H.) in Trondheim, and by the research nurse (A.C.) in Groningen.<sup>23</sup> To quantify extracranial injuries, a modified extracranial score ( $ISS_e$ ) was calculated based on the total ISS score minus the squared Abbreviated Injury Scale Head score.<sup>24</sup>

### Head CT

Most head CTs were acquired with a Siemens Somatom Sensation 64-row scanner (Siemens AG, Erlangen, Germany) in both Trondheim and Groningen. CT examinations were performed as standardized care at ED admission and during follow-up if needed. Both the first and the worst CT examinations were reviewed for the current study by a consultant or resident in neurosurgery or radiology (I.H.S., K.G.M., S.F.D., or J.X.) in Trondheim and in Groningen by a neurologist (J.V.N. or B.J.).

At both centers, head CT characteristics of the worst CT examination were categorized into: any intracranial finding, SDH, EDH, traumatic subarachnoid hemorrhage (tSAH), including intraventricular hemorrhage, punctate hemorrhage  $<2$  mm, contusion (single or multiple lesions), fracture (base, skull, and impression fractures merged), midline shift  $>$  or  $\leq 5$  mm, and degree of compression of basal cisterns (normal, compressed, or absent). In addition, Rotterdam CT score (best score 1, worst score 6) using the worst scan was computed by consultant or resident in neurosurgery or radiology (I.H.S., K.G.M., S.F.D., or J.X.) in Trondheim and in Groningen by a consultant in physical medicine and rehabilitation (C.E.) based on the CT variables.<sup>25</sup> This score is based on midline shift, compression of basal cisterns, tSAH, or intraventricular hemorrhage and EDH.

### Outcome

Length of stay was defined as time from the first hospital ED admission to discharge from the level I trauma center. Case-fatality rate was defined as the percentage of patients who died from the head injury during the hospital stay. Discharge destinations were home (with or without outpatients' rehabilitation services), other clinical departments (including psychiatry), other hospitals, rehabilitation center (including rehabilitation at hospital, rehabilitation in private institutions or municipal rehabilitation), or nursing home (including sheltered housing 24 hours a day).

Functional outcome was assessed at 12 months after the injury using the structured interview for the GOSE.<sup>26-29</sup> The GOSE score in survivors was assessed based on the functional outcome from the injury as a whole and not specifically the brain injury. Outcome was assessed by phone (most of the patients at Trondheim) or in-person interview (most of the patients at Groningen) with the patients and relatives or caregivers. Both phone and in-person interview for the GOSE assessment have been validated and good agreement has been found, especially after standardizing procedures and training.<sup>26,30,31</sup> The outcome assessors were not blinded for clinical information.

The GOSE score was dichotomized into being moderately disabled or worse (GOSE score  $\leq 6$ ) versus good recovery (GOSE score 7–8), and into being severely disabled or worse (GOSE score  $\leq 4$ ) versus moderate or no disability (GOSE score 5–8).

### Statistical Analysis

The statistical analyses were conducted with IBM SPSS Statistics version 22 (IBM Corp., Armonk, New York, USA), STATA/SE

**Table 1.** All Patients: Clinical Characteristics and Injury-Related Variables

Variable	Total	Trondheim	Groningen	P Value
No. of patients, %	395	190 (48)	205 (52)	
Age, years				<b>0.002</b>
Median (range, IQR)	46 (16–97, 25–63)	51 (16–97, 28–67)	39 (16–88, 23–60)	
Mean $\pm$ SD	46 $\pm$ 21	50 $\pm$ 22	43 $\pm$ 39	
Male/female sex, <i>n</i> (%)	280/115 (71/29)	123/67 (65/35)	157/48 (77/23)	<b>0.010</b>
Preinjury disability, <i>n</i> (%)*	92 (24)	49 (26)	43 (22)	0.305
Cause of injury, <i>n</i> (%)*				<b>0.006</b>
Traffic accidents	174 (45)	71 (39)	103 (50)	
Fall	166 (43)	94 (51)	72 (35)	
Other	49 (13)	19 (10)	30 (15)	
GCS score, <i>n</i> (%)				0.921
13	171 (43)	84 (44)	87 (42)	
12	79 (20)	39 (21)	40 (20)	
11	44 (11)	22 (12)	22 (11)	
10	40 (10)	19 (10)	21 (10)	
9	61 (15)	26 (14)	35 (17)	
Alcohol intoxication, <i>n</i> (%)†	109 (28)	48 (26)	61 (31)	0.277
Intoxicated, S-ethanol known	51 (13)	41 (22)	10 (5)	
Intoxicated, S-ethanol unknown	58 (15)	7 (4)	51 (26)	
Transferred from other hospitals, <i>n</i> (%)	64 (16)	41 (22)	23 (11)	<b>0.005</b>
Secondary event, <i>n</i> (%)‡	50 (13)	22 (12)	28 (14)	0.609
Unilateral pupillary dilation, <i>n</i> (%)§	42 (11)	15 (8)	27 (13)	0.086
Median ISS score (IQR)	17 (9–25)	20 (13–25)	17 (8–24)	<b>&lt;0.001</b>
Median ISS <sub>e</sub> score (IQR)	4 (0–9)	1 (0–8)	4 (1–9)	<b>&lt;0.001</b>
Intracranial findings, <i>n</i> (%)	320 (81)	173 (91)	147 (72)	<b>&lt;0.001</b>
SDH, <i>n</i> (%)	136 (34)	88 (46)	48 (23)	<b>&lt;0.001</b>
EDH, <i>n</i> (%)	62 (16)	29 (15)	33 (16)	0.820
tSAH, <i>n</i> (%)	195 (49)	106 (56)	89 (43)	<b>0.014</b>
Punctate hemorrhage, <i>n</i> (%)	108 (27)	48 (25)	60 (29)	0.372
Contusion(s), <i>n</i> (%)	181 (46)	107 (56)	74 (36)	<b>&lt;0.001</b>
Cranial fracture, <i>n</i> (%)	174 (44)	87 (46)	87 (42)	0.503
Midline shift >5 mm, <i>n</i> (%)	66 (17)	34 (18)	32 (16)	0.543
Basal cisterns, <i>n</i> (%)				0.374
Normal	300 (76)	150 (79)	150 (73)	
Compressed	84 (21)	36 (19)	48 (23)	
Absent	11 (3)	4 (2)	7 (3)	
Median Rotterdam CT score (IQR)	3 (2–3)	3 (2–3)	2 (2–3)	<b>0.030</b>
Admitted to ICU, <i>n</i> (%)	279 (71)	167 (88)	112 (55)	<b>&lt;0.001</b>
Median days ICU LOS (IQR)	4 (2–8)	3 (1–7)	5 (2–12)	<b>&lt;0.001</b>
Intubated, <i>n</i> (%)	164 (42)	87 (46)	77 (38)	0.097

Continues

Table 1. Continued

Variable	Total	Trondheim	Groningen	P Value
Median days on ventilator (IQR)¶	4 (1–10)	2 (1–9)	6 (1–11)	<b>0.045</b>
ICP monitoring, n (%)#	66 (17)	44 (23)	22 (11)	<b>0.001</b>
Evacuation of mass lesion, n (%)	60 (15)	39 (21)	21 (10)	<b>0.004</b>
Median days LOS (range, IQR)**	8 (0–142, 3–16)	6 (1–94, 4–12)	10 (0–142, 3–20)	0.055
In-hospital case-fatality rate, n (%)	13 (3)	6 (3)	7 (3)	0.886
Discharge destination, n (%)††				<b>&lt;0.001</b>
Home	280 (48)	57 (31)	123 (63)	
Other hospital	93 (25)	68 (37)	25 (27)	
Rehabilitation	74 (20)	37 (20)	37 (19)	
Nursing home	24 (6)	14 (8)	10 (5)	
Other departments	7 (2)	6 (3)	1 (1)	

Significant P values are marked in bold.

IQR, interquartile range; SD, standard deviation; GCS, Glasgow Coma Scale; ISS, Injury Severity Score; ISS<sub>e</sub>, modified ISS score for extracranial injuries; SDH, subdural hematoma; EDH, epidural hematoma; tSAH, traumatic subarachnoid hemorrhage; CT, computed tomography; ICU, Intensive care unit; LOS, length of hospital stay; ICP, intracranial pressure.

Number of patients in analysis due to lacking data: \*n = 389, †n = 388, ‡n = 380, §n = 394.

¶Only patients treated in ICU included, n = 275.

¶¶Only ventilated patients included, n = 163.

#Reasons for ICP monitoring were high-risk brain injury (79%) or monitoring of the brain injury in sedated patients (21%).

\*\*Only surviving patients included, n = 380.

††Only surviving patients included, n = 378.

version 13 (StataCorp LLC, College Station, Texas, USA), and the R statistical package.<sup>32</sup> Demographic and injury characteristics are presented as percentages, median with interquartile range (IQR), or mean with standard deviation. Between-group differences were analyzed with the Mann–Whitney U test for variables with non-normal distributions and for ordinal variables. The  $\chi^2$  test or Fisher exact test was used for comparison of proportions. Two-sided P values of  $< 0.05$  were considered statistically significant.

A common model was generated based on the data from Trondheim and Groningen combined and one for each of the Trondheim and Groningen datasets. We included 9 clinical and 8 CT variables commonly used in previous outcome prediction studies. We chose to use the individual CT characteristics rather than the Rotterdam CT score, because they are more easily interpreted in a clinical context. Age, GCS score, and ISS<sub>e</sub> score were analyzed as continuous variables, and the remaining clinical and CT variables were binary. In addition, center was included as a

Table 2. All Patients: 12 Months' Outcome

GOSSE Score at 12 Months	Total	Trondheim	Groningen	P Value
No. of patients	358	160 (45)	198 (55)	
GOSSE 1 (Death), n (%)	20 (6)	10 (6)	10 (5)	
GOSSE 3 (Severe disability, lower), n (%)	16 (4)	12 (8)	4 (2)	
GOSSE 4 (Severe disability, upper), n (%)	14 (4)	4 (3)	10 (5)	
GOSSE 5 (Moderate disability, lower), n (%)	45 (13)	17 (11)	28 (14)	
GOSSE 6 (Moderate disability, upper), n (%)	63 (18)	24 (15)	39 (20)	
GOSSE 7 (Good recovery, lower), n (%)	82 (23)	25 (16)	57 (29)	
GOSSE 8 (Good recovery, upper), n (%)	118 (33)	68 (43)	50 (25)	
GOSSE score $\leq 6$ (%), n (%)	158 (44)	67 (42)	91 (46)	0.439
GOSSE score $\leq 4$ (%), n (%)	50 (14)	26 (16)	24 (12)	0.263

GOSSE, Glasgow Outcome Scale Extended.



**Table 3.** Patients Included in the Prediction Analysis: Clinical Characteristics, Injury-Related Variables, and 12 Months' Outcome

Variable	Total	Trondheim	Groningen	P Value
No. of patients (%)	329	147 (45)	182 (55)	
Age, years				0.138
Median (range, IQR)	45 (16–97, 24–62)	48 (16–97, 25–63)	39 (16–88, 23–61)	
Mean $\pm$ SD	45 $\pm$ 21	47 $\pm$ 21	43 $\pm$ 21	
Male/female sex, <i>n</i> (%)	232/97 (71/30)	93/54 (63/37)	139/43 (76/24)	<b>0.010</b>
Preinjury disability, <i>n</i> (%)	69 (21)	30 (20)	39 (21)	0.821
Cause of injury, <i>n</i> (%)				0.094
Traffic accident	155 (47)	61 (42)	94 (52)	
Fall	133 (40)	69 (47)	64 (35)	
Other	41 (13)	17 (12)	24 (13)	
Median GCS score (IQR)	12 (10–13)	12 (10–13)	12 (10–13)	0.399
Alcohol intoxication, <i>n</i> (%)	93 (28)	39 (27)	54 (30)	0.530
Secondary event, <i>n</i> (%)	39 (12)	12 (8)	27 (15)	0.063
Unilateral pupillary dilation, <i>n</i> (%)	34 (10)	11 (8)	23 (13)	0.127
Median ISS <sub>e</sub> (IQR)	4 (1–9)	4 (0–8)	4 (1–9)	<b>0.001</b>
SDH, <i>n</i> (%)	109 (33)	65 (44)	44 (24)	<b>&lt;0.001</b>
EDH, <i>n</i> (%)	56 (17)	25 (17)	31 (17)	0.995
tSAH, <i>n</i> (%)	164 (50)	82 (56)	82 (45)	0.053
Punctate hemorrhage, <i>n</i> (%)	98 (30)	40 (27)	58 (32)	0.358
Contusion(s), <i>n</i> (%)	148 (45)	81 (55)	67 (37)	<b>0.001</b>
Cranial fracture, <i>n</i> (%)	146 (44)	66 (45)	80 (44)	0.864
Midline shift >5 mm, <i>n</i> (%)	65 (17)	27 (18)	29 (16)	0.559
Basal cisterns, <i>n</i> (%)				0.446
Normal	246 (75)	114 (78)	132 (73)	
Compressed	73 (22)	30 (20)	43 (24)	
Absent	10 (3)	3 (2)	7 (4)	
Median Rotterdam CT score (IQR)	3 (2–3)	3 (2–3)	2 (2–3)	0.068
GOSE score $\leq$ 6, <i>n</i> (%)	147 (45)	62 (42)	85 (47)	0.412

Significant *P* values are marked in bold.IQR, interquartile range; SD, standard deviation; GCS, Glasgow Coma Scale; ISS<sub>e</sub>, modified ISS score for extracranial injuries; SDH, subdural hematoma; EDH, epidural hematoma; tSAH, traumatic subarachnoid hemorrhage; CT, computed tomography; GOSE, Glasgow Outcome Scale Extended.

binary covariate (fixed effect) in the common model, in effect accounting for within-center dependencies. Missing data were handled by listwise deletion (13 patients in Trondheim: 5 cause of injury and 8 secondary event [Figure 1A] and 16 patients in Groningen: 5 preinjury disability, 5 day-of-injury alcohol intoxication, 5 secondary event, 1 pupillary dilation [Figure 1B]). A total of 329 patients were included in the analysis.

The models were fitted by penalized logistic regression using the lasso (least absolute shrinkage and selection operator) method as implemented in the R package glmnet.<sup>33</sup> This method shrinks the values of the regression coefficients to obtain less extreme values,

as a means towards improving the external validity of the model. For variables with low predictive value, the coefficients could be shrunk to zero, and the variables thus left out of the final model. The degree of shrinkage was determined by 10-fold cross-validation. In effect, the method performs simultaneous estimation of the coefficients and variable selection. It should be noted that the lasso method focuses on the overall fit rather than statistical significance of individual predictors. Consequently, predictors with a *P* value > 0.05 could still be included in the final model.

The uncertainty in the estimated coefficients from the lasso was assessed by bootstrapping the penalized regression procedure

**Table 4.** Common Model Selected by Lasso

Variable	Coefficient	OR (95% CI)	P Value
Intercept	0.84		
Age	0.02	1.02 (1.02–1.04)	<b>&lt;0.001</b>
Female	0.09	1.09 (0.76–2.26)	0.324
Preinjury disability	0.40	1.50 (1.02–3.40)	<b>0.043</b>
Traffic accident	0.06	1.06 (0.71–2.08)	0.480
GCS score	−0.24	0.65 (0.65–0.90)	<b>0.001</b>
Alcohol intoxication	−0.83	0.44 (0.20–0.65)	<b>0.001</b>
Secondary event	0.62	1.86 (1.05–5.14)	<b>0.037</b>
Pupillary dilation	0.37	1.44 (0.85–4.45)	0.113
ISS <sub>e</sub>	0	1 (0.97–1.05)	0.628
SDH	0.62	1.86 (1.31–4.14)	<b>0.004</b>
EDH	0.12	1.13 (0.63–2.53)	0.520
tSAH	0.26	1.30 (0.91–2.63)	0.107
Punctate hemorrhage	0	1 (0.67–2.05)	0.574
Contusion	0	1 (0.51–1.57)	0.710
Cranial fracture	0.17	1.19 (0.71–2.26)	0.433
Midline shift >5 mm	0.13	1 (0.67–3.34)	0.323
Basal cisterns compressed/absent	0.18	1.20 (0.68–2.63)	0.404
Center	0.23	1.26 (0.92–2.54)	0.102

Significant *P* values are marked in bold.

Models selected by lasso show the estimated shrunk regression coefficients for the combined Trondheim and Groningen data. A coefficient of 0 means that the variable was not included in the model and values different from 0, was included. OR was the odds for GOSE score ≤6 versus odds for GOSE score >6.

OR, odds ratio; CI, confidence interval; GCS, Glasgow Coma Scale; ISS<sub>e</sub>, Injury Severity Scale extracranial; SDH, subdural hematoma; EDH, epidural hematoma; tSAH, traumatic subarachnoid hemorrhage.

using 1000 bootstrap samples. A bootstrap sample is generated by resampling with replacement from the original data set. The penalized regression procedure, including the selection of the degree of shrinkage, was run for each bootstrap sample. The uncertainty was illustrated for each of the variables by the proportion of the 1000 bootstrap samples that gave a value of zero for that coefficient. Proportions closer to zero indicate greater probability for the variable to be included in the model and contribute to the outcome prediction. In addition, *P* values for the regression coefficients were calculated using the de-sparsified lasso method implemented in the R package hdi.<sup>34</sup> This method takes the shrinkage and variable selection into account. *P* values < 0.05 were considered statistically significant. Because the statistical analysis was aimed at obtaining the best predictor model, the statistical significance for the regression coefficients was not of major importance, and no formal adjustment for multiple testing was included.

The model fit was assessed by the Hosmer–Lemeshow goodness-of-fit test (HL test), for which a *P* value less than 0.05 indicates a poor fit. The Nagelkerke pseudo-*R*<sup>2</sup> was also calculated

for the models. The area under the curve (AUC) of the receiver operating characteristics curves was used to assess discrimination. The 95% confidence intervals (CIs) for the AUCs and the *P* values for comparing AUCs were calculated by bootstrapping using 10,000 bootstrap samples. To calculate the performance measures for external validation, the Groningen outcomes were predicted based on the model fitted in the Trondheim data, and vice versa.

### Ethics

In Trondheim, the Regional Committee for Medical Research Ethics (2013/1977) approved the study. Written consent was obtained from the surviving patients or their next of kin if the patient was incapacitated. In Groningen, the study was approved by the local Medical Ethical Committee, and informed consent was waived as only de-identified clinical data were registered.

### RESULTS

#### All Patients with Moderate TBI

In the total study population of 395 patients, the median age was 46 years, 71% of the patients were male, and 24% reported preinjury disability (Table 1). Falls and traffic accidents were the main causes of injury. Intracranial traumatic lesions on head CT were seen in 81% of the patients, 71% were treated in an ICU (median 4 days (IQR = 2–8 days), and median days on ventilator were 4 (IQR = 1–10). The in-hospital case-fatality rate was 3%. At 12-month follow-up, 56% had good recovery (GOSE score 7 and 8), 31% moderate disability (GOSE score 5 and 6), 8% severe disability (GOSE score 3 and 4), and 6% had died from their head injury (GOSE score 1). Hence, 44% were moderately disabled or worse (GOSE score ≤6), whereas 14% were severely disabled or worse (GOSE score ≤4) (Table 2).

The GCS score was 13 in 43% of the patients. In general, the GCS 9–12 group were significantly different from the GCS 13 group, with a greater proportion of intracranial lesions (73% vs. 88%, *P* ≤ 0.001), treatment in ICU (56% vs. 82%, *P* ≤ 0.001), and GOSE score ≤6 at follow-up (35% vs. 51%, *P* = 0.003). Still, the GCS 13 group had high prevalence of lesions on CT, more than 50% were treated in the ICU, and approximately one third did not achieve a good recovery.


#### Differences Between the Two Centers

In Trondheim, the patients were older (median 51 years vs. 39 years), the proportion of women was greater, and more injuries were caused by falls than in Groningen. The patients in Trondheim had more often been transferred from other hospitals and traumatic intracranial lesions were more often present on head CT (91% vs. 72%). They had greater ISS scores, but ISS<sub>e</sub> was lower. The patients in Trondheim were more often admitted to the ICU and more often had ICP monitoring (Table 1). More patients were discharged directly home in Groningen than in Trondheim (63% vs. 31%), whereas discharge to other hospitals was more common in Trondheim (37% vs. 27%). Outcome at 12 months was not significantly different between the centers.

#### Prediction of Outcome

Characteristics of the patients included in the outcome prediction analysis are presented in Table 3. The combined dataset of both





Trondheim and Groningen was used to develop a model for prediction of a GOSE score  $\leq 6$ , including center as an additional categorical variable (Table 4, Figure 2A). Of the variables selected as predictors for GOSE score  $\leq 6$ , older age ( $P < 0.001$ ), lower GCS score ( $P = 0.001$ ), no day-of-injury alcohol intoxication ( $P = 0.001$ ), presence of SDH ( $P = 0.004$ ), occurrence of a secondary event ( $P = 0.037$ ), and preinjury disability ( $P = 0.043$ ) were significant associated with outcome. The HL test for the common model indicated a good model fit ( $P = 0.143$ ). The Nagelkerke pseudo- $R^2$  was 0.34, and AUC for the common prognostic model was 0.80 (95% CI 0.75–0.85) (Figure 3).


In the Trondheim cohort, older age ( $P < 0.001$ ), pupillary dilation ( $P = 0.005$ ), the presence of a SDH ( $P = 0.012$ ), and preinjury disability ( $P = 0.049$ ) were significantly associated with a GOSE score  $\leq 6$  (Figure 2B). The Nagelkerke pseudo- $R^2$  was 0.43, the HL test indicated a good model fit ( $P = 0.325$ ), and the AUC from the internal validation was 0.85 (95% CI 0.78–0.91) (Figure 3). In Groningen, a lower GCS score ( $P = 0.001$ ), no day-of-injury alcohol intoxication ( $P = 0.002$ ), older age ( $P = 0.007$ ), and the presence of a SDH ( $P = 0.030$ ) were significantly associated with the outcome (Figure 2C). The Nagelkerke

pseudo- $R^2$  was 0.31, the HL test indicated a good model fit ( $P = 0.270$ ), and the AUC from the internal validation was 0.79 (95% CI 0.72–0.85) (Figure 3).

When the Trondheim model was tested in the Groningen data, the AUC value was 0.75 (95% CI 0.68–0.82), and the  $P$  value of the HL-test was 0.038. When the Groningen model was tested in the Trondheim data, the AUC value was 0.76 (95% CI 0.67–0.83) and the  $P$  value for the HL-test was 0.362.

## DISCUSSION

In this follow-up study performed exclusively in patients with moderate TBI from 2 European level 1 trauma centers, approximately three quarters of the patients had intracranial findings on head CT, and many needed intensive care treatments. Few patients died or experienced severe disability. Still, 44% of the patients did not achieve good recovery at 12 months. Older age, lower GCS score, no day-of-injury alcohol intoxication, SDH, occurrence of secondary event, and preinjury disability were predictors for GOSE score  $\leq 6$  in a model that was constructed from the combined dataset.



We found that 81% had intracranial traumatic lesions on head CT, which is somewhat greater than previously reported.<sup>1-3,10</sup> The lowest frequency was reported in the oldest study and may be explained by lower detection rates with older scanners. The most frequent intracranial findings in the present study were tSAH and contusions, in line with other studies on patients with moderate TBI.<sup>1,10</sup> Acute SDH was reported less frequently, only in 34%, whereas this lesion type is more common in severe TBI.<sup>1,35</sup>

Further, the patients with moderate TBI often were treated in an ICU and many underwent neurosurgical interventions. ICP monitoring was performed in several patients as might not be expected in patients with moderate TBI. In most of these cases, the patients had a high-risk brain injury, whereas some patients needed sedation for other reasons, and the ICP measurement was implemented to monitor the evolution of the brain injury. Despite all these indicators of significant brain injury, the in-hospital case-fatality rate was low (3%), in line with other studies on patients with moderate TBI (0.9%–8%).<sup>2,3,5,10</sup> This finding lends validity to the debated GCS score as a clinically useful tool for classification of injury severity in the acute setting.<sup>36</sup> The current study also demonstrated that the patients with GCS score 13 suffered significant injuries as they had a high rate of intracranial CT findings and 35% had a disability at 12 month follow-up. Indeed, it is debated whether patients with GCS score 13 should

be classified as mild or moderate TBI.<sup>8,37</sup> Based on the current results in a large sample, the clinical characteristics of patients with GCS score 13 provide evidence for these individuals as belonging to the moderate TBI rather than mild TBI category. This classification scheme is in line with the Head Injury Severity Scale and also has been used in previous studies.<sup>1,2,9</sup>

The significant cohort effects between the cohorts from Trondheim and Groningen underline the necessity for multicenter studies. Ongoing international multisite studies, like the TRACK-TBI and the CENTER-TBI, are important in this respect. The approach of comparative effectiveness research in these studies will hopefully increase our understanding of the provided care and measured outcomes of patients across centers.

A GOSE score  $\leq 4$ , i.e., death or severe disability, was observed in only 14% in the present study. This was in accordance with 2 Italian studies reporting GOSE score  $\leq 4$  at 6 months, in 15% and 26%, respectively.<sup>2,5</sup> Hence, supported by results from our study, we argue that the existing models like CRASH and IMPACT, which are designed to predict such poor outcomes, have low relevance for patients with moderate TBI. In contrast, a GOSE score of  $\leq 6$ , i.e., worse than good recovery, was observed in 44% of the patients in the present study, quite similar to the 55% reported in a previous Dutch multicenter study from 5 level I trauma centers in which patients with GCS score 13 also were included.<sup>1</sup> The high number of patients with moderate TBI with persistent functional problems call for routine follow-up in these patients, who are often discharged to their homes.<sup>20</sup>

### Prediction of Outcome

This is the first study specifically to develop a prognostic model in a sample of only patients with moderate TBI. Moreover, we applied a cut-off at moderate disability or worse (GOSE score  $\leq 6$ ), unlike previous studies, which have developed and validated models that predict severe disability or death (GOSE score of  $\leq 4$ ). In the model fitted based on the combined dataset, older age, lower GCS score, no day-of-injury alcohol intoxication, SDH, occurrence of a secondary event, and preinjury disability were significant predictors of GOSE score  $\leq 6$ .

Age is a well-known prognostic factor for outcome in patients with TBI.<sup>15,16,38</sup> Studies on patients with severe TBI show greater case-fatality rate for elderly patients,<sup>39,40</sup> as well as worse long-term outcome.<sup>40,41</sup> The present study clearly shows that this is also the case for patients with moderate TBI. Also in accordance with findings in severe TBI, the GCS score, occurrence of a secondary event, and SDH were related to worse outcome.<sup>11,15,16,42</sup>

The variable “pre-injury disability” was associated with worse outcome in the entire sample and the Trondheim cohort but not in the Groningen cohort. This variable might have been defined differently and hence subject to between-center variations. Still, this result indicates that adding variables describing preinjury health may increase the prognostic performance of a model, as also has been shown for mild TBI.<sup>43,44</sup> Therefore, we suggest that future prognostic studies of patients with moderate TBI should explore the impact of pre- and comorbidity.

A more surprising finding was that not being influenced by alcohol was associated with worse outcome, most prominent in the Groningen cohort. However, positive serum ethanol has previously also been associated with a better outcome.<sup>45,46</sup> One

explanation of this finding could be that the depressant effects of alcohol on the central nervous system was falsely ascribed to the head injury.<sup>19</sup> If so, patients with only mild TBI could be included in cohorts of moderate and severe TBI due to falsely low GCS score and present with a good recovery at follow-up. In contrast, alcohol has also been hypothesized to have a neuroprotective effect previous studies.<sup>47-48</sup> Regardless, influence of alcohol is an example of clinical information that probably should be systematically collected at admission and controlled for in future studies of prognosis.

The common outcome prediction model performed adequately with an AUC value of 0.80, but the discriminative ability needs to be proven by external validation. Yet, we believe that the model presented here comprises variables that represent important risk factors for disability after moderate TBI. The modest Nagelkerke pseudo- $R^2$  of the model indicates that the outcome after moderate TBI may depend also on factors that are not measured in regular moderate-severe TBI cohort studies. Future studies could address this shortcoming by collection of a broader set of variables embracing a biopsychosocial understanding of the TBI patient in line with the insight from the field of mild TBI.<sup>44,49</sup>

In the external validations of the 2 separate models, the models performed similarly, with AUC values of 0.76 (Trondheim) and 0.75 (Groningen), compared with 0.85 and 0.79, respectively, for the internal validations. Thus, the AUC values indicate that the models show an acceptable strength of discrimination between moderate disability or worse in contrast to good recovery.<sup>50</sup> However, to arrive at a predictive model with sufficient discriminative ability to be used in clinical practice, further studies are needed. We believe that an important challenge regarding prognostication in moderate TBI is to identify and incorporate the best set of factors that may influence outcome in the individual patients. Especially important is it to extend future studies to middle- and low-income countries, and the International Initiative for TBI Research (InTBIR) is promising in this respect.<sup>51</sup>

### Strengths and Limitations

A strength of this study is the large number of prospectively registered patients with moderate TBI. Since we specifically studied patients with moderate TBI, we could choose the level of dichotomization of the GOSE that we believe is the most relevant, according to the baseline risk of patients with moderate TBI. Moreover, the chosen statistical method, penalized regression using the lasso method, is an important strength. It performs simultaneous estimation of the coefficients and the variable

selection, determining the best combination of variables with reduced risk of overfitting of the models.

One limitation of the study was that inclusion and data collection was planned and completed separately in the 2 centers. Moreover, both cohorts comprise only patients who have been treated at neurosurgical referral centers, and the study results may not apply to patients who are treated in general hospitals. We doubt, however, that this has caused a large bias in this study, because both hospitals also serve as general hospitals, and because most patients with moderate TBI are referred to a neurosurgical center in both countries. Another limitation is the wide time-span of data-collection, which can have impact on performance of the model, but is difficult to avoid when a large sample is needed. Finally, it is important to bear in mind that this study describes patients treated in 2 high-income countries, and our conclusions may therefore not be valid in middle and low-income countries.

### CONCLUSIONS

In this prospective study from 2 European centers, a high proportion of the patients with moderate TBI had characteristics of significant brain injury and needed advanced hospital care. Therefore, it is important to secure appropriate acute care. Even if few patients died, a high proportion (44%) did not experience a good recovery, which substantiates the need of appropriate follow-up for patients with moderate TBI.

Older age, lower GCS score, no day-of-injury alcohol intoxication, SDH, occurrence of secondary event, and preinjury disability were predictors for GOSE score  $\leq 6$ . Future studies should incorporate an even broader set of variables, which can hopefully increase the predictive power of a prognostic model. We believe that this study can serve as a first step in future development of valid prognostic models for patients with moderate TBI.

### ACKNOWLEDGMENTS

The authors acknowledge the following contributors in Trondheim: consultants in radiology Ingrid Haavde Strand and Susan Frances Deane and resident in radiology Jian Xu for classification of the CT scans; consultant in neurosurgery Sozaburo Hara for performing some of the ISS calculations; and research nurses Beate M. Voll, Oddrun Sandrød, Stine Borgen Lund, and occupational therapist Otto Aarhaug for the management of the patient database and performing the GOSE interviews. In Groningen, the authors acknowledge Annemiek Coers for management of the patient database.

### REFERENCES

- Andriessen TM, Horn J, Franschman G, van der Naalt J, Haitsma I, Jacobs B, et al. Epidemiology, severity classification, and outcome of moderate and severe traumatic brain injury: a prospective multicenter study. *J Neurotrauma*. 2011;28:2019-2031.
- Fabbri A, Servadei F, Marchesini G, Stein SC, Vandelli A. Early predictors of unfavourable outcome in subjects with moderate head injury in the emergency department. *J Neurol Neurosurg Psychiatry*. 2008;79:567-573.
- Fearnside M, McDougall P. Moderate head injury: a system of neurotrauma care. *Aust N Z J Surg*. 1998;68:58-64.
- Lund SB, Gjeilo KH, Moen KG, Schirmer-Mikalsen K, Skandsen T, Vik A. Moderate traumatic brain injury, acute phase course and deviations in physiological variables: an observational study. *Scand J Trauma Resusc Emerg Med*. 2016;24:77.
- Compagnone C, d'Avella D, Servadei F, Angileri FF, Brambilla G, Conti C, et al. Patients with moderate head injury: a prospective multicenter study of 315 patients. *Neurosurgery*. 2009;64:690-696.
- Rimel RW, Giordani B, Barth JT, Jane JA. Moderate head injury: completing the clinical spectrum of brain trauma. *Neurosurgery*. 1982;11:344-351.
- Teasdale G, Jennett B. Assessment of coma and impaired consciousness. A practical scale. *Lancet*. 1974;2:81-84.

8. Maas AI, Stocchetti N, Bullock R. Moderate and severe traumatic brain injury in adults. *Lancet Neurol*. 2008;7:728-741.
9. Stein SC, Spettell C. The Head Injury Severity Scale (HISS): a practical classification of closed-head injury. *Brain Inj*. 1995;9:437-444.
10. Stein SC, Ross SE. Moderate head injury: a guide to initial management. *J Neurosurg*. 1992;77:562-564.
11. Steyerberg EW, Mushkudiani N, Perel P, Butcher I, Lu J, McHugh GS, et al. Predicting outcome after traumatic brain injury: development and international validation of prognostic scores based on admission characteristics. *PLoS Med*. 2008;5:e165.
12. Tagliaferri F, Compagnone C, Korsic M, Servadei F, Kraus J. A systematic review of brain injury epidemiology in Europe. *Acta Neurochir (Wien)*. 2006;148:255-268.
13. Peeters W, van den Brande R, Polinder S, Brazinova A, Steyerberg EW, Lingsma HF, et al. Epidemiology of traumatic brain injury in Europe. *Acta Neurochir (Wien)*. 2015;157:1683-1696.
14. Husson EC, Ribbers GM, Willemse-van Son AH, Verhagen AP, Stam HJ. Prognosis of six-month functioning after moderate to severe traumatic brain injury: a systematic review of prospective cohort studies. *J Rehabil Med*. 2010;42:425-436.
15. MRC CRASH Trial Collaborators, Perel P, Arango M, Clayton T, Edwards P, Komolafe E, et al. Predicting outcome after traumatic brain injury: practical prognostic models based on large cohort of international patients. *BMJ*. 2008;336:425-429.
16. Jacobs B, Beems T, van der Vliet TM, van Vugt AB, Hoedemaekers C, Horn J, et al. Outcome prediction in moderate and severe traumatic brain injury: a focus on computed tomography variables. *Neurocrit Care*. 2013;19:79-89.
17. Maas AIR, Menon DK, Adelson PD, Andelic N, Bell MJ, Belli A, et al. Traumatic brain injury: integrated approaches to improve prevention, clinical care, and research. *Lancet Neurol*. 2017;16:987-1048.
18. Skandsen T, Kvistad KA, Solheim O, Strand IH, Folvik M, Vik A. Prevalence and impact of diffuse axonal injury in patients with moderate and severe head injury: a cohort study of early magnetic resonance imaging findings and 1-year outcome. *J Neurosurg*. 2010;113:556-563.
19. Rundhaug NP, Moen KG, Skandsen T, Schirmer-Mikalsen K, Lund SB, Hara S, et al. Moderate and severe traumatic brain injury: effect of blood alcohol concentration on Glasgow Coma Scale score and relation to computed tomography findings. *J Neurosurg*. 2015;122:211-218.
20. de Koning ME, Spikman JM, Coers A, Schonherr MC, van der Naalt J. Pathways of care the first year after moderate and severe traumatic brain injury-discharge destinations and outpatient follow-up. *Brain Inj*. 2015;29:423-429.
21. Puljula J, Savola O, Tuomivaara V, Pribula J, Hillbom M. Weekday distribution of head traumas in patients admitted to the emergency department of a city hospital: effects of age, gender and drinking pattern. *Alcohol*. 2007;42:474-479.
22. Mahler SA, Pattani S, Standifer J, Caldito G, Conrad SA, Arnold TC. Clinical sobriety assessment by emergency physicians in blunt trauma patients with acute alcohol exposure. *J Emerg Med*. 2010;39:685-690.
23. Baker SP, O'Neill B, Haddon W Jr, Long WB. The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma*. 1974;14:187-196.
24. Gennarelli TA, Wodzin E. AIS 2005: a contemporary injury scale. *Injury*. 2006;37:1083-1091.
25. Maas AI, Hukkelhoven CW, Marshall LF, Steyerberg EW. Prediction of outcome in traumatic brain injury with computed tomographic characteristics: a comparison between the computed tomographic classification and combinations of computed tomographic predictors. *Neurosurgery*. 2005;57:1173-1182.
26. Wilson JT, Pettigrew LE, Teasdale GM. Structured interviews for the Glasgow Outcome Scale and the extended Glasgow Outcome Scale: guidelines for their use. *J Neurotrauma*. 1998;15:573-585.
27. Jennett B, Snoek J, Bond MR, Brooks N. Disability after severe head injury: observations on the use of the Glasgow Outcome Scale. *J Neurol Neurosurg Psychiatry*. 1981;44:285-293.
28. Wilde EA, Whiteneck GG, Bogner J, Bushnik T, Cifu DX, Dikmen S, et al. Recommendations for the use of common outcome measures in traumatic brain injury research. *Arch Phys Med Rehabil*. 2010;91:1650-1660.
29. Pettigrew LE, Wilson JT, Teasdale GM. Reliability of ratings on the Glasgow Outcome Scales from in-person and telephone structured interviews. *J Head Trauma Rehabil*. 2003;18:252-258.
30. Wilson JT, Edwards P, Fiddes H, Stewart E, Teasdale GM. Reliability of postal questionnaires for the Glasgow Outcome Scale. *J Neurotrauma*. 2002;19:999-1005.
31. Wilson JT, Slieker FJ, Legrand V, Murray G, Stocchetti N, Maas AI. Observer variation in the assessment of outcome in traumatic brain injury: experience from a multicenter, international randomized clinical trial. *Neurosurgery*. 2007;61:123-128.
32. R Development Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Available at: <https://www.R-project.org/>; 2016. Accessed June 1, 2017.
33. Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via Coordinate Descent. *J Stat Softw*. 2010;33:1-22.
34. Dezeure R, Buhlmann P, Meier L, Meinshausen N. High-dimensional inference: confidence intervals, p-values and R-Software hdi. *Stat Sci*. 2015;30:533-558.
35. Gomez PA, Castano-Leon AM, de-la-Cruz J, Lora D, Lagares A. Trends in epidemiological and clinical characteristics in severe traumatic brain injury: analysis of the past 25 years of a single centre data base. *Neurocirugia (Astur)*. 2014;25:199-210.
36. Teasdale G, Maas A, Lecky F, Manley G, Stocchetti N, Murray G. The Glasgow Coma Scale at 40 years: standing the test of time. *Lancet Neurol*. 2014;13:844-854.
37. Mena JH, Sanchez AI, Rubiano AM, Peitzman AB, Sperry JL, Gutierrez MI, et al. Effect of the modified Glasgow Coma Scale score criteria for mild traumatic brain injury on mortality prediction: comparing classic and modified Glasgow Coma Scale score model scores of 13. *J Trauma*. 2011;71:1185-1192.
38. Maas AI, Marmarou A, Murray GD, Teasdale SG, Steyerberg EW. Prognosis and clinical trial design in traumatic brain injury: the IMPACT study. *J Neurotrauma*. 2007;24:232-238.
39. Utomo WK, Gabbe BJ, Simpson PM, Cameron PA. Predictors of in-hospital mortality and 6-month functional outcomes in older adults after moderate to severe traumatic brain injury. *Injury*. 2009;40:973-977.
40. Roe C, Skandsen T, Manskow U, Ader T, Anke A. Mortality and one-year functional outcome in elderly and very old patients with severe traumatic brain injuries: observed and predicted. *Behav Neurol*. 2015;2015:845-849.
41. de Guise E, LeBlanc J, Dagher J, Tinawi S, Lamoureux J, Marcoux J, et al. Traumatic brain injury in the elderly: a level 1 trauma centre study. *Brain Inj*. 2015;29:558-564.
42. Roozenbeek B, Lingsma HF, Lecky FE, Lu J, Weir J, Butcher I, et al. Prediction of outcome after moderate and severe traumatic brain injury: external validation of the International Mission on Prognosis and Analysis of Clinical Trials (IMPACT) and Corticoid Randomisation After Significant Head injury (CRASH) prognostic models. *Crit Care Med*. 2012;40:1609-1617.
43. Cnossen MC, Polinder S, Andriessen TM, van der Naalt J, Haitsma I, Horn J, et al. Causes and consequences of treatment variation in moderate and severe traumatic brain injury: a multicenter study. *Crit Care Med*. 2017;45:660-669.
44. van der Naalt J, Timmerman ME, de Koning ME, van der Horn HJ, Scheenen ME, Jacobs B, et al. Early predictors of outcome after mild traumatic brain injury (UPFRONT): an observational cohort study. *Lancet Neurol*. 2017;16:532-540.

45. Jacobs B, Beems T, Stulemeijer M, van Vugt AB, van der Vliet TM, Borm GF, et al. Outcome prediction in mild traumatic brain injury: age and clinical variables are stronger predictors than CT abnormalities. *J Neurotrauma*. 2010;27:655-668.
46. Scheenen ME, de Koning ME, van der Horn HJ, Roks G, Yilmaz T, van der Naalt J, et al. Acute alcohol intoxication in patients with mild traumatic brain injury: characteristics, recovery, and outcome. *J Neurotrauma*. 2016;33:339-345.
47. Raj R, Skrifvars MB, Kivisaari R, Hernesniemi J, Lappalainen J, Siironen J. Acute alcohol intoxication and long-term outcome in patients with traumatic brain injury. *J Neurotrauma*. 2015;32: 95-100.
48. Alexander S, Kerr ME, Yonas H, Marion DW. The effects of admission alcohol level on cerebral blood flow and outcomes after severe traumatic brain injury. *J Neurotrauma*. 2004;21: 575-583.
49. Cnossen MC, Winkler EA, Yue JK, Okonkwo DO, Valadka A, Steyerberg EW, et al. Development of a prediction model for post-concussive symptoms following mild traumatic brain injury: a TRACK-TBI Pilot Study. *J Neurotrauma*. 2017;34:2396-2409.
50. Hosmer DW, Lemeshow S, Sturdivant RX. *Applied Logistic Regression*. 3rd ed. Hoboken, NJ: John Wiley & Sons; 2013.
51. Tosetti P, Hicks RR, Theriault E, Phillips A, Koroshetz W, Draghia-Akli R, et al. Toward an international initiative for traumatic brain

injury research. *J Neurotrauma*. 2013;30: 1211-1222.

*Conflict of interest statement:* This work was supported by funding from the Liaison Committee between the Central Norway Regional Health Authority (RHA) and the Norwegian University of Science and Technology (NTNU).

Received 23 December 2017; accepted 24 March 2018

Citation: *World Neurosurg.* (2018) 114:e1199-e1210.

<https://doi.org/10.1016/j.wneu.2018.03.176>

Journal homepage: [www.WORLDNEUROSURGERY.org](http://www.WORLDNEUROSURGERY.org)

Available online: [www.sciencedirect.com](http://www.sciencedirect.com)

1878-8750/© 2018 The Author(s). Published by Elsevier Inc.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).